

## LA-UR-19-30005

Approved for public release; distribution is unlimited.

Title: Dark Matter Searches with HAWC

Author(s): Lundeen, Joseph Andrew

Intended for: International Cosmic Ray Conference, 2019-07-24 (Madison, Wisconsin, United States)

Issued: 2019-10-02

---

**Disclaimer:**

Los Alamos National Laboratory, an affirmative action/equal opportunity employer, is operated by Triad National Security, LLC for the National Nuclear Security Administration of U.S. Department of Energy under contract 89233218CNA000001. By approving this article, the publisher recognizes that the U.S. Government retains nonexclusive, royalty-free license to publish or reproduce the published form of this contribution, or to allow others to do so, for U.S. Government purposes. Los Alamos National Laboratory requests that the publisher identify this article as work performed under the auspices of the U.S. Department of Energy. Los Alamos National Laboratory strongly supports academic freedom and a researcher's right to publish; as an institution, however, the Laboratory does not endorse the viewpoint of a publication or guarantee its technical correctness.



# Dark Matter Searches with HAWC


Joe Lundeen for the HAWC Collaboration

July, 2019



# Continuous Spectra

- WIMP dark matter can annihilate or decay to standard model particles
  - Interaction products then produce photons
- Energy spectrum characterized by hard cutoff at DM mass
- Search for gamma-ray excesses with characteristic shape originating from known DM halos
- Can constrain velocity-weighted cross section or decay lifetime
- Emphasis on multi-TeV mass dark matter




$$\frac{d\Phi}{dE_{\text{annihilation}}} = \frac{J}{8\pi} \frac{\langle\sigma v\rangle}{M^2} \frac{dN(M, \text{channel})}{dE}$$

$$J = \int \int \rho_{dm}^2(l, \Omega) dl d\Omega$$

$$\frac{d\Phi}{dE_{\text{decay}}} = \frac{D}{4\pi} \frac{1}{\tau M} \frac{dN(M, \text{channel})}{dE}$$

$$D = \int \int \rho_{dm}(l, \Omega) dl d\Omega$$






# HAWC Detector

**Citlaltepetl**  
Pico de Orizaba  
5160m a.s.l.

- **22,000 m<sup>2</sup> air shower array**
- **300 Water Cherenkov detectors (WCD)**
- **200,000 liters of purified water per WCD**
- **4 sensors (photo-multiplier tubes) per WCD**
- **Completed March 2015**
- **Near-continuous duty cycle**
- **Field of view within 2/3 of sky**
- **Ideal for all-sky surveys**



*Large Millimeter Telescope*




**HAWC**  
4100 m a.s.l.

**Tliltepetl**  
Sierra Negra  
4582m a.s.l.




# HAWC Properties and Advantages

- Wide simultaneous field of view ( $\sim 2$  sr)
  - Sensitive to highly-extended sources
  - Direct integration for background estimation
- Observation of  $\sim 2/3$  of sky every day
  - Ability to survey for new sources
  - Can search for DM in multiple regions simultaneously  $\rightarrow$  combined searches
- Archival data
- Sensitivity is declination-dependent
  - Due to atmospheric attenuation of showers
  - Better sensitivity to sources that transit overhead



Abeysekara et. al. *Astrophys.J.* 843 (2017) no.1, 40

# Dark Matter Search Targets





# Dwarf Galaxies

- Excellent candidates for DM searches
- Relatively sparse star population
  - No known normal-matter production mechanism for high-energy gamma-rays
  - Very little astrophysical background
- Continuous duty cycle:
  - Can easily perform combined limits
  - Can add additional limits as more are discovered

- Two Classes:
  - Dwarf Spheroidal
    - 15 candidates
    - See next slide
  - Dwarf Irregular
    - 31 candidates
    - See next talk by Sergio Hernandez



# Limits from Dwarf Spheroidals



Note: The J-factor of Triangulum II is not well known. Limits are reported with and without Tri II.  
See: A. Albert et al., *Astrophys. J.* 853 (2018) 154.

# Joint Limits


- Joint search of the dwarf spheroidals with multiple experiments
  - HAWC, HESS, MAGIC, VERITAS, Fermi-LAT
  - Complete coverage of all multi-GeV through multi-TeV dark matter masses
  - First ever analysis of its kind
- See talk from Monday by Louise Oakes and proceedings



+



+



# Extended Targets

- M31 galaxy, Virgo Cluster, Galactic halo
- Wide field of view needed
- Allows for full treatment of morphology
- Background estimation
  - Need “off” regions sufficiently far from source to avoid signal contamination
  - Wide field of view and continuous duty-cycle allows for simultaneous observation of “on” and “off” regions
- Need to consider systematics from spatial profile
- Particularly well-suited for setting decay lower-limits

# Density Profile Uncertainty

- Behavior of dark matter density not well constrained towards center of large halos
- J-factors and D-factors typically have large systematic from density profile
- Signal boosts from theorized substructure contribution


## Einasto Profile (Cuspy)

$$\rho(r) = \rho_s e^{\frac{-2}{\alpha} [(r/r_s)^\alpha - 1]}$$

## Burkert Profile (Cored)

$$\rho(r) = \frac{\rho_s}{(1 + r/r_s)(1 + (r/r_s)^2)}$$


Dark Matter Radial Profiles







# M31

- Closest galaxy (besides the Milky Way)
  - Combined with high dark matter content, gives large expected flux
- Also highly extended
  - Requires treatment of morphology
  - Considered different density profiles
- Substructure
  - Different models of substructure content considered
  - Results shown for median model
- Yields strong decay limits




See: Albert et. al. *Search for Dark Matter Gamma-ray Emission from the Andromeda Galaxy with the High-Altitude Water Cherenkov Observatory*

# Decay Limits from M31





# The Virgo Cluster

- Highly extended
  - $\sim 10 \times 10$  degrees
  - Morphology consists of two distinct peaks
- Different models of substructure contribution
  - High, median, and low substructure content models
  - Only results from median case shown here
- High dark matter content
- Nicely compliments constraints from other experiments
- See poster presentation by Tolga Yapici for details




# Decay Limits from Virgo




# The Galactic Halo

- Closest large halo → large expected flux
- Largest flux expected towards the Galactic center, however:
  - Large systematic from unconstrained density profile
  - Possible contamination from visible-matter sources
- HAWC field of view enables observation of larger regions further from the center
  - Mitigates effect of density profile
  - Avoids contamination from sources in Galactic plane
- Previous analysis in Fermi Bubble region (see: HAWC Collaboration, A. U. Abeysekara et al., JCAP 1802 (2018) 049.)
- Current analysis using even wider region (see poster and proceedings by Joe Lundeen)



# Gamma-ray Lines


- Direct annihilation of dark matter to gamma-rays
- Manifests as a delta function in energy spectrum
- “Smoking gun” for dark matter
  - Only mechanism that can produce this shape at TeV scale
  - Location of line immediately reveals the dark matter mass.
- New energy estimation techniques allow HAWC to search for this feature





# First Limits on Gamma-ray Lines

- Performed combined upper limit using dwarf spheroidal galaxies
- Most constraining limits above  $\sim 20$  TeV
- Nicely compliment searches by IACTs



# Summary

- HAWC's wide field of view and continuous duty cycle make it ideal for surveys and extended source analysis
- Can easily perform combined searches
  - Improves sensitivity of constraints
  - Currently extending to combinations with other experiments
- Sensitive to extended sources: yield strong decay limits
- Now able to search for gamma-ray lines
  - Energy estimators also allow searches at higher masses
- More results coming from Galactic halo





# Backup Slides



# Likelihood Fitting

- Use log likelihood ratio used to identify signals
  - N=observed counts in each bin
  - B=expected background
  - S=expected excess counts from dark matter spectrum
  - Treat counts in each bin as Poisson-distributed
  - Calculate likelihoods for both null (background) hypothesis and dark matter hypothesis
- Wilks' theorem
  - TS is chi-squared distributed
  - Allows us to treat  $\sigma$  as significance for 1 degree of freedom
- 95% confidence level
  - Fit such that  $TS=TS_{\text{max}}-2.71$

$$L_0 = \frac{e^{-B} B^N}{N!}$$


$$L = \frac{e^{-(B+S)} (B+S)^N}{N!}$$

$$TS = -2 \ln \left( \frac{L_0}{L} \right)$$


$$\sigma = \text{sign}(\phi) \sqrt{TS}$$

# Forward Folding

- Due to energy resolution, HAWC does not bin events by energy
- Binned by fraction of PMTs triggered (fhit)
- fhit spectrum converted back to energy spectrum
  - Use simulation of detector response to energy spectra
  - Map observed nhit spectrum back to E-spectrum that best reproduces it in simulation
  - Use maximum likelihood



Forward folding



# Direct Integration

- Majority of HAWC events are charged cosmic rays
  - Roughly isotropic across the sky
  - Some are removed through quality cuts, but still dominate post-cut data
- Background estimated through direct integration
  - Average event rate within 2 hours (30 degrees) of a source
  - Events are background-dominated → average rate is approximately background rate
- R. Atkins, et al. *Astrophys.J.* 595 (2003) 803-811
- See also: D. Fiorino, Thesis.

# Density Profile Uncertainty

- Behavior of dark matter density not well constrained towards center of large halos
- J-factors and D-factors typically have large systematic from density profile


## Einasto Profile (Cuspy)

$$\rho(r) = \rho_s e^{\frac{-2}{\alpha} [(r/r_s)^\alpha - 1]}$$

## Burkert Profile (Cored)

$$\rho(r) = \frac{\rho_s}{(1 + r/r_s)(1 + (r/r_s)^2)}$$

Dark Matter Radial Profiles




# The Galactic Halo

- Closest large halo → high expected dark matter signal
- HAWC can observe highly extended region of the halo
- Requires very careful treatment of background
  - Extends across entire sky
  - If dark matter does produce gamma rays, all sky regions will contain some contribution from Galactic halo
  - No true “Off” region for background estimation
  - Need to take into account signal contamination in background
- Highly sensitive to density profile

# The Galactic Halo


- Closest large halo
- Requires very careful treatment of background
  - Extends across entire sky
  - If dark matter does produce gamma rays, all sky regions will contain some contribution from Galactic halo
  - No true “Off” region for background estimation
  - Need to take into account signal contamination in background
- Highly sensitive to density profile

# Decay Limits from the Galactic Halo



Obtained these limits using just a fairly small part of the halo.


Limits will improve with the more rigorous search (see poster by Joe Lundeen)





# Galactic Halo Sensitivity

- Obtained by combining estimates of dark matter sensitivity with modeled density profile
- Map of point source “expected limits”: show which pixels are expected to be most constraining
  - Most sensitive pixels are combined into full ROI
  - Remaining sky is used as “off” region of background estimation



Galactic center is at edge of HAWC field of view  
Limits are mostly driven by points overhead